

# revisi Bessel

*by Rizky Azis*

---

**Submission date:** 18-Nov-2023 10:57AM (UTC+0700)

**Submission ID:** 2231891612

**File name:** Cauchy\_Rizky\_Aziz\_S\_Autosaved\_1\_4.pdf (577.08K)

**Word count:** 2499

**Character count:** 11338



## 5 The Boundedness of Generalized Fractional Integral Operators on Small Morrey Spaces

Corina Karim<sup>1</sup>, Rizky Aziz Syaifudin<sup>1</sup>, Marjono<sup>1</sup>, Hairur Rahman<sup>2</sup>

<sup>1</sup>Department of Mathematic, Brawijaya University, Indonesia

<sup>2</sup>Department of Mathematic, Islamic University of Maulana Malik Ibrahim, Indonesia

Email: [co\\_mathub@ub.ac.id](mailto:co_mathub@ub.ac.id), [rizkyaziz@student.ub.ac.id](mailto:rizkyaziz@student.ub.ac.id), [marjono@ub.ac.id](mailto:marjono@ub.ac.id),  
[hairur@mat.uin-malang.ac.id](mailto:hairur@mat.uin-malang.ac.id)

### 19 ABSTRACT

The small Morrey space is the set of locally Lebesgue integrable functions with norm defined supremum over radius of ball  $r \in (0,1)$ . This paper aims to prove the boundedness properties of the generality of fractional integral operators in small Morrey spaces using Hedberg-type inequality. The first, in this paper will be discuss to prove Hedberg-type inequality on small Morrey spaces using dyadic decomposition, Hölder inequality, and doubling condition. Furthermore, by using the inequality, it can be proven that the boundedness of generalized fractional integral operators on small Morrey spaces.

**Keywords:** Fractional Integral Operators; Hedberg-type inequality; small Morrey spaces;

### INTRODUCTION

15  
Morrey space was first introduced by C. B. Morrey in 1938. Morrey [1] found a solution space of elliptic differential equations called Morrey space. In the development of research, Morrey space develops into several kinds of spaces such as small Morrey space. The fractional integral operator is one type of singular integral operator. Research on fractional integral operators, first started by Hardy and Littlewood. Hardy and Littlewood [2] proved the boundedness of fractional integral operators on Lebesgue spaces using dyadic decomposition. Furthermore, the research was continued by Adams [3] and Chiarenza and Frasca [4] who proved the boundedness of fractional integral operators on Morrey space using dyadic decomposition.

1  
Nakai [5] generalized the space of Adams and Chiarenza and Frasca. Nakai [5] proved the boundedness property of fractional integral operators on the generalized Morrey spaces using dyadic decomposition. The result of Nakai research is also called Spanne inequality. Gunawan and Eridani [6] extended the results of Adams and Chiarenza and Frasca. Karim and Apriliani [7] continued their research on the Spanne inequality of the  $p = q$  case.

Sawano [8] extended Morrey space into a small Morrey space for  $r \in (0,1)$ . In Sawano's research [8], it also gives the properties of inclusions among small Morrey spaces. In Sawano's research, it also gives the properties of inclusions among small Morrey spaces. Based on the inclusion properties, Karim et. al. [9] research further discussed the inclusion of Lebesgue spaces, Morrey spaces, and small Morrey spaces. Karim, et.al. [9] has proved the boundedness of fractional integral operator in small Morrey spaces using Hedberg Inequality.

## METHODS

The method used in writing this research article is a literature study of several related articles and books. In some of the literature, the definition and limitation properties of fractional integral operators have been discussed and the limitation properties and definition of Morrey space have also been discussed. Therefore, by using the definitions and properties that have been discussed in the previous literature study, it can be used to prove the boundedness properties in this article. The first, we will discuss about Lebesgue spaces and Morrey spaces from (see [10], [11], [12], [13], [14], and [15]).

### Definition 1 (Lebesgue Spaces)

Let  $(X, \mathcal{A}, \mu)$  is measurable spaces and  $1 \leq p < \infty$ . The Lebesgue spaces  $L^p(X)$  are consists all equivalence class of function  $f : X \rightarrow \mathbb{R}$  such that

$$\|f\|_{L^p} := \left( \int_X |f|^p dx \right)^{\frac{1}{p}} < \infty.$$

### Definition 2 (Local Lebesgue Spaces)

Let  $(X, \mathcal{A}, \mu)$  is measurable spaces. The Lebesgue spaces  $L^p_{loc}(X)$  are consists all equivalence class of function  $f : X \rightarrow \mathbb{R}$  such that for every compact subset  $S \subset X$ ,

$$\int_S |f|^p dx < \infty.$$

### Theorem 1 (Hölder Inequality)

If  $f \in L^p(\mathbb{R}^n)$  dan  $g \in L^q(\mathbb{R}^n)$  with  $p > 1$ , then  $\|fg\|_{L^1} \leq \|f\|_{L^p} \|g\|_{L^q}$ .

*Proof.* The proof of this theorem can be seen in [10].

### Definition 2 (Fractional Integral Operator)

Let  $0 < \beta < n$ . The integral fractional operator  $I_\beta f(x)$  is defined by

$$I_\beta f(x) := \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\beta}} dy$$

for every  $f \in L^p_{loc}(\mathbb{R}^n)$  with  $1 \leq p < \infty$  and for every  $x \in \mathbb{R}^n$ .

### Theorem 2 (Hedberg-type inequality)

Let  $1 \leq q < p < \infty$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\beta}{n}$ , then for every  $f \in L^p(\mathbb{R}^n)$ ,

$$|I_\beta f(x)| \leq C_{p,q} (Mf(x))^{1-\frac{\beta p}{n}} \|f(x)\|_{L^p}^{\frac{\beta p}{n}}.$$

*Proof.* Suppose  $1 \leq q < p < \infty$ , take any  $f \in L^p(\mathbb{R}^n)$  and open ball  $O(x, r)$ . From definition integral fractional operator, write

$$I_\beta f(x) := I_1 f(x) + I_2 f(x),$$

with

$$I_1 f(x) := \int_{O(x,r)} \frac{f(y)}{|x-y|^{n-\beta}} dy,$$

and

$$I_2 f(x) := \int_{O(x,r)^c} \frac{f(y)}{|x-y|^{n-\beta}} dy.$$

Using dyadic decomposition, partition of the open ball  $O(x,r)$  is  $2^k r \leq |x-y| < 2^{k+1} r$  with  $-\infty < k \leq -1$  such that

$$\begin{aligned} |I_1 f(x)| &\leq C \sum_{k=-\infty}^{-1} \int_{2^k r \leq |x-y| < 2^{k+1} r} \frac{f(y)}{|x-y|^{n-\beta}} dy \\ &\leq C r^\beta (Mf(x)). \end{aligned}$$

Partition of the complement of open ball  $O(x,r)^c$ , we obtain  $2^k r \leq |x-y| < 2^{k+1} r$  with  $0 \leq k < \infty$  such that

$$\begin{aligned} |I_2 f(x)| &\leq C \sum_{k=0}^{\infty} \int_{2^k r \leq |x-y| < 2^{k+1} r} \frac{f(y)}{|x-y|^{n-\beta}} dy \\ &\leq C r^{\beta-\frac{n}{p}} \|f(x)\|_{L^p}. \end{aligned}$$

Choose  $r := \left( \frac{\|f(x)\|_{L^p}}{Mf(x)} \right)^{\frac{p}{n}}$ , such that we obtain

$$\begin{aligned} |I_\beta f(x)| &\leq |I_1 f(x)| + |I_2 f(x)| \\ &\leq C \left( r^\beta (Mf(x)) + r^{\beta-\frac{n}{p}} \|f(x)\|_{L^p} \right) \\ &\leq C (Mf(x))^{1-\frac{\beta p}{n}} \|f(x)\|_{L^p}^{\frac{\beta p}{n}}. \end{aligned}$$

So, Hedberg-type Inequality is holds.

### Definition 3 (Doubling Condition)

In defining the operators  $I_\psi$ , the function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$  satisfy the doubling condition if holds

1. If  $\frac{1}{2} \leq \frac{r}{s} \leq 2$  then  $\frac{1}{C_1} \leq \frac{\psi(r)}{\psi(s)} \leq C_1$  for every  $r, s > 0$ .
2. For  $r > 0$ ,  $\int_r^\infty \frac{\psi(t)}{t} dt$ .

### Definition 4 (Morrey Spaces)

Let  $0 \leq q \leq p < \infty$  and measurable spaces  $X = \mathbb{R}^n$ . The Morrey spaces  $\mathcal{M}_q^p(\mathbb{R}^n)$  consists of function  $f \in L_{loc}^p(\mathbb{R}^n)$  with

$$\|f\|_{\mathcal{M}_q^p} := \sup_{x \in \mathbb{R}^n, r > 0} |O(x, r)|^{\frac{1}{q} - \frac{1}{p}} \left( \int_{O(x, r)} |f(x)|^p dx \right)^{\frac{1}{p}},$$

is finite.

**Definition 5** (Small Morrey Spaces)

Let  $0 \leq q \leq p < \infty$  and measurable spaces  $X = \mathbb{R}^n$ . The small Morrey spaces  $m_q^p(\mathbb{R}^n)$  consists of function  $f \in L_{loc}^p(\mathbb{R}^n)$  with

$$\|f\|_{m_q^p} := \sup_{x \in \mathbb{R}^n, r \in (0, 1)} |O(x, r)|^{\frac{1}{q} - \frac{1}{p}} \left( \int_{O(x, r)} |f(x)|^p dx \right)^{\frac{1}{p}},$$

is finite.

**RESULTS AND DISCUSSIONS**

In this section, we will discuss the boundedness properties of the fractional integral operator. However, we will first prove Hedberg-type inequality in small Morrey spaces. Eridani [11] generalized the fractional integral operator. Given function  $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  that satisfies the doubling condition. The generalized fractional integral operator is defined by

$$I_\psi f(x) := \int_{\mathbb{R}^n} \frac{\psi(|x - y|)}{|x - y|^n} f(y) dy,$$

for every  $f \in L_{loc}^p(\mathbb{R}^n)$  with  $1 \leq p < \infty$ . If  $\psi(|x - y|) := |x - y|^\beta$  with  $0 < \alpha < n$  then  $I_\psi f(x) = I_\beta f(x)$  [14].

Since  $\rho$  function satisfies the doubling condition, such that for every  $k \in \mathbb{Z}$  and for every  $r > 0$  with  $2^k r \leq |x - y| < 2^{k+1} r$  holds

1. If  $\frac{1}{2} \leq \frac{|x-y|}{2^{k+1}r} \leq 2$  then

$$\frac{1}{C} \leq \frac{\psi(|x - y|)}{\psi(2^{k+1}r)} \leq C.$$

As a result, we obtain

$$\frac{\psi(2^{k+1}r)}{C} \leq v(|x - y|) \leq C\psi(2^{k+1}r). \quad (1)$$

2. Let  $t := |x - y|$ . The inequality (1) can be expressed as

$$\frac{\psi(2^{k+1}r)}{C} \leq \psi(t) \leq C\psi(2^{k+1}r). \quad (2)$$

The next, from inequality (2), we have

$$\begin{aligned} \int_{2^k r}^{2^{k+1} r} \frac{\psi(2^{k+1}r)}{C 2^{k+1}r} dt &\leq \int_{2^k r}^{2^{k+1} r} \frac{\psi(t)}{t} dt \leq \int_{2^k r}^{2^{k+1} r} \frac{C\psi(2^{k+1}r)}{2^k r} dt \\ \frac{\psi(2^{k+1}r)}{2C} &\leq \int_{2^k r}^{2^{k+1} r} \frac{\psi(t)}{t} dt \leq C\psi(2^{k+1}r). \end{aligned} \quad (3)$$

**Theorem 3** (Hedberg-type Inequality)

Let function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$  satisfies doubling condition. If  $1 \leq p < q < \infty$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\beta}{n}$ , then

$$|I_\psi f(x)| \leq C_{p,q} |Mf(x)|^{1-\frac{\beta q}{n}} \|f(x)\|_{m_q^p}^{\frac{\beta q}{n}},$$

for every  $f \in m_q^p(\mathbb{R}^n)$ , with  $Mf(x)$  is maximal Hardy-Littlewood operator.

*Proof.* Take any  $f \neq 0 \in m_q^p(\mathbb{R}^n)$  and let  $1 \leq p < q < \infty$ . Defined a function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$ , is  $\psi(t) := t^\beta$  with  $0 < \beta < n$  satisfies doubling condition such that  $\psi(t)$  satisfies inequality (1) and (3). Write

$$I_\psi f(x) := I_1 f(x) + I_2 f(x),$$

With

$$I_1 f(x) := \int_{O(x,r)} \frac{\psi(|x-y|)}{|x-y|^n} f(y) dy,$$

and

$$I_2 f(x) := \int_{O(x,r)^c} \frac{\psi(|x-y|)}{|x-y|^n} f(y) dy.$$

For open ball  $O(x,r)$ , can be partitioned  $2^k r \leq |x-y| < 2^{k+1} r$  for every  $-\infty < k \leq -1$  such that we obtain

$$\begin{aligned} |I_1 f(x)| &\leq \sum_{k=-\infty}^{-1} \int_{2^k r \leq |x-y| < 2^{k+1} r} \frac{\psi(|x-y|)}{|x-y|^n} |f(y)| dy \\ &\leq C \sum_{k=-\infty}^{-1} \psi(2^{k+1} r) \frac{1}{(2^{k+1} r)^n} \left( \int_{|x-y| < 2^{k+1} r} |f(y)| dy \right) \\ &= C \sum_{k=-\infty}^{-1} (2^{k+1} r)^\beta \left( \frac{1}{(2^{k+1} r)^n} \int_{|x-y| < 2^{k+1} r} |f(y)| dy \right) \\ &\leq C \sum_{k=-\infty}^{-1} (2^{k+1} r)^\beta \left( \sup_{O(x, 2^{k+1} r)} \frac{1}{(2^{k+1} r)^n} \int_{|x-y| < 2^{k+1} r} |f(y)| dy \right) \\ &= C \sum_{k=-\infty}^{-1} (2^{k+1} r)^\beta Mf(x). \end{aligned}$$

Because the series  $\sum_{k=-\infty}^{-1} 2^{\beta(k+1)}$  converges to one for  $k < 0$ , then inequality can be expressed by

$$|I_1 f(x)| \leq C r^\beta Mf(x).$$

For  $O(x,r)^c$ , the ball  $O(x,r)^c$  can be partitioned  $2^k r \leq |x-y| < 2^{k+1} r$  with  $0 \leq k < \infty$ . Observe that

$$\begin{aligned} |I_2 f(x)| &\leq \sum_{k=0}^{\infty} \int_{2^k r \leq |x-y| < 2^{k+1} r} \frac{\psi(|x-y|)}{|x-y|^n} |f(y)| dy \\ &\leq C \sum_{k=0}^{\infty} \frac{\psi(2^k r)}{(2^k r)^n} \int_{|x-y| < 2^{k+1} r} \frac{\psi(|x-y|)}{|x-y|^n} |f(y)| dy, \end{aligned}$$

with constant  $C > 0$  and using Hölder inequality we have

$$\frac{1}{(2^k r)^n} \int_{|x-y| < 2^{k+1} r} |f(y)| dy \leq \left( \frac{1}{(2^k r)^n} \int_{|x-y| < 2^{k+1} r} |f(y)|^p dy \right)^{\frac{1}{p}}$$

$$\leq (2^k r)^{-\frac{n}{q}} \|f\|_{m_q^p}.$$

Because  $\psi(t) := t^\beta$ , consequently can be expressed as

$$\begin{aligned} |I_2 f(x)| &\leq C \sum_{k=0}^{\infty} \frac{\psi(2^k r)}{(2^k r)^n} \int_{|x-y| < 2^{k+1} r} \frac{\psi(|x-y|)}{|x-y|^n} |f(y)| dy \\ &\leq C \sum_{k=0}^{\infty} (2^k r)^\beta (2^k r)^{-\frac{n}{q}} \|f\|_{m_q^p} \\ &\leq C r^{\beta-\frac{n}{q}} \sum_{k=0}^{\infty} (2^k)^{\beta-\frac{n}{q}} \|f\|_{m_q^p} \end{aligned}$$

and because  $0 < \beta < n$  then  $\beta - n < \beta - \frac{n}{q} < 0$  for  $1 < p < q < \infty$  such that the series  $\sum_{k=0}^{\infty} (2^k)^{\beta-\frac{n}{q}}$  converges to one. The result is

$$|I_2 f(x)| \leq C r^{\beta-\frac{n}{q}} \|f\|_{m_q^p}.$$

Next, using triangle inequality we obtain

$$\begin{aligned} |I_\psi f(x)| &\leq |I_1 f(x)| + |I_2 f(x)| \\ &\leq C r^\beta Mf(x) + C r^{\beta-\frac{n}{q}} \|f\|_{m_q^p} \\ &\leq C \left( r^\beta Mf(x) + r^{\beta-\frac{n}{q}} \|f\|_{m_q^p} \right). \end{aligned}$$

Choose  $r := \left( \frac{\|f\|_{m_q^p}}{Mf(x)} \right)^{\frac{q}{n}}$ , such that

$$|I_\psi f(x)| \leq C \left( r^\beta Mf(x) + r^{\beta-\frac{n}{q}} \|f\|_{m_q^p} \right) = C (Mf(x))^{1-\frac{\beta q}{n}} \|f\|_{m_q^p}^{\frac{\beta q}{n}}.$$

So, Hedberg-type inequality on small Morrey spaces for fractional integral operator is proved.

The following presents the limitation theorem of the generalized fractional integral operator in small Morrey spaces.

#### Theorem 4

Let positive function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$  satisfies doubling condition and  $1 < p_1 < q_1 < \frac{n}{\beta}$  with  $0 < \beta < n$ . If  $\frac{1}{q_2} = \frac{1}{q_1} - \frac{\beta}{n}$  and  $\frac{1}{p_2} = \frac{1}{p_1} \left( 1 - \frac{\beta q_1}{n} \right)$ , then exist constant  $C_{p_1, q_1} > 0$  such that for every  $f \in m_{q_1}^{p_1}(\mathbb{R}^n)$  holds

$$\|I_\psi f\|_{m_{q_2}^{p_2}} \leq C_{p_1, q_1} \|f\|_{m_{q_1}^{p_1}}.$$

*Proof.* Let  $1 < p_1 < q_1 < \frac{n}{\beta}$  with  $0 < \beta < n$ . Defined function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$ , is  $\psi(t) := t^\beta$ . Take any function  $f \neq 0 \in m_{q_1}^{p_1}(\mathbb{R}^n)$ . According to Hedberg-type Inequality from Theorem 3, holds

$$|I_\psi f(x)| \leq C_{p, q} |Mf(x)|^{1-\frac{\beta q}{n}} |f(x)|_{m_q^p}^{\frac{\beta q}{n}}.$$

Defined

$$\frac{1}{q_2} = \frac{1}{q_1} - \frac{\beta}{n},$$

and

$$\frac{1}{p_2} = \frac{1}{p_1} \left(1 - \frac{\beta q_1}{n}\right).$$

Therefore, it is obtained

$$\begin{aligned} |I_\psi f(x)| &\leq C(Mf(x))^{1-\frac{\beta n}{q_1}} \|f\|_{m_{q_1}^{p_1}}^{\frac{\beta n}{q_1}} \\ &= C(Mf(x))^{\frac{p_1}{p_2}} \|f\|_{m_{q_1}^{p_1}}^{1-\frac{p_1}{p_2}}, \end{aligned}$$

Next, both segments are raised by  $p_2$  and integrated over  $O(x, r)$  such that

$$\begin{aligned} \int_{O(x,r)} |I_\psi f(x)|^{p_2} dx &\leq C^{p_2} (Mf(x))^{p_1} \|f\|_{m_{q_1}^{p_1}}^{p_2-p_1} \\ &= C^{p_2} \|f\|_{m_{q_1}^{p_1}}^{p_2-p_1} \int_{B(x,r)} (Mf(x))^{p_1} dx \\ &\leq C^{p_2} \|f\|_{m_{q_1}^{p_1}}^{p_2-p_1} \sup_{x \in \mathbb{R}^n, r \in (0,1)} |O(x, r)|^{\frac{1}{q_1} - \frac{1}{p_1}} \int_{O(x,r)} (Mf(x))^{p_1} dx \\ &= C^{p_2} \|f\|_{m_{q_1}^{p_1}}^{p_2-p_1} \|Mf\|_{m_{q_1}^{p_1}}^{p_1} \\ &\leq C^{p_2} \|f\|_{m_{q_1}^{p_1}}^{p_2-p_1} \|f\|_{m_{q_1}^{p_1}}^{p_1} \\ &\leq C^{p_2} \|f\|_{m_{q_1}^{p_1}}^{p_2}. \end{aligned}$$

Rank both segments by  $\frac{1}{p_2}$  then multiply by  $|O(x, r)|^{\frac{1}{q_2} - \frac{1}{p_2}}$  and take the supremum over radius  $r \in (0,1)$  of both segments such that we get

$$\|I_\psi f\|_{m_{q_2}^{p_2}} \leq C_{p_1, q_1} \|f\|_{m_{q_1}^{p_1}}.$$

Thus, it is proven that  $I_\psi f(x)$  is finite in the small Morrey spaces.

## CONCLUSIONS

The generalization of the fractional integral operators by defining the positive function  $\psi : \mathbb{R} \rightarrow \mathbb{R}^+$  that satisfies the doubling condition. The proof of the boundedness properties of the generalized fractional integral operator in small Morrey space uses the Hedberg-type inequality. Therefore, in proving it, it is necessary to prove first that the Hedberg-type inequality applies in small Morrey space and applies to the generalized fractional integral operator.

## REFERENCES

- [1] Morrey, C.B. 1938. On the Solutions of Quasi-linear Elliptic Partial Differential Equations. *Trans. Amer. Math. Soc.* 43: 126-166.

- [2] Hardy, G.H. dan Littlewood, J.E. 1927. Some Properties of Fractional Integrals. *Math. Zeit.* 27: 565-605.
- [3] Adams, D. 1975. A Note on Riesz Potentials. *Duke Maths.* 42: 765-778.
- [4] Chiarenza, F. dan Frasca, M. 1987. Morrey Spaces and Hardy-Littlewood Maximal Function. *Rend. Mat.* 7: 273-279.
- [5] Nakai, E. 1994. Hardy-Littlewood Maximal Operator, Singular Integral Operator, and Riesz Potentials on Generalized Morrey Spaces. *Math. Nachr.* 166: 95-103
- [6] Gunawan, H. and Eridani. 2009. Fractional integrals and generalized Olsen inequalities. *Kyungpook Math.* 49(1):31-39.
- [7] Karim, C., dan Apriliani, E. 2008. *Variasi Ketaksamaan Spanne*. Prosiding Seminar Nasional Matematika IV. Surabaya: 13 Desember 2008.
- [8] Sawano, Y., Fazio, G. D., and Maryani, S. 2020. *Introduction and Applications to Integral Operators and PDE's*. CRC Press.
- [9] Karim, C., Syaifudin, R. A., Marjono, and Rahman, H. 2023. The boundedness of fractional integral operators on small Morrey spaces. Atlantis Press Proceeding, Brawijaya International Conference.
- [10] Sawano, Y., Fazio, G. D., and Maryani, S. 2020. *Introduction and Applications to Integral Operators and PDE's*. CRC Press.
- [11] Bartle, R. G. dan D.R. Sherbert. 1995. *The Elements of Integration and Lebesgue Measure*. John Willey & Sons. New York.
- [12] Sawano, Y. 2018. A Thought on Generalized Morrey Spaces. *J. Indonesian. Math. Soc.* 3(25): 210-281.
- [13] Eridani. 2002. On the Boundedness of a Generalized Fractional Integral on Generalized Morrey Spaces. *Tamkang Journal of Mathematics.* 4(33): 335-340.
- [14] Eridani. 2002. On the Boundedness of A Generalized Fractional Integral on Generalized Morrey Spaces. *Tamkang Journal of Mathematics.* 4(33): 335-340.
- [15] Castillo, R. E. dan H. Rafeiro. 2016. *An Introductory Course in Lebesgue Spaces*. Springer. Swiss.

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

10%

INTERNET SOURCES

16%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

[1library.net](http://1library.net)

Internet Source

3%

2

Michael Ruzhansky, Durvudkhan Suragan. "Hardy Inequalities on Homogeneous Groups", Springer Science and Business Media LLC, 2019

Publication

2%

3

Arun Sharma, Dinesh K. Kumar. "Chapter 20 Moments and Wavelets for Classification of Human Gestures Represented by Spatio-Temporal Templates", Springer Science and Business Media LLC, 2004

Publication

2%

4

[etheses.uin-malang.ac.id](http://etheses.uin-malang.ac.id)

Internet Source

1%

5

Hendra Gunawan, Denny Ivanal Hakim, Yoshihiro Sawano, Idha Sihwaningrum. "Weak Type Inequalities for Some Integral Operators on Generalized Nonhomogeneous Morrey Spaces", Journal of Function Spaces and Applications, 2013

1%

|    |                                                                                                                                                                                                                                              |      |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 6  | Michael Ruzhansky, Durvudkhan Suragan, Nurgissa Yessirkegenov. "Hardy-Littlewood, Bessel-Riesz, and fractional integral operators in anisotropic Morrey and Campanato spaces", Fractional Calculus and Applied Analysis, 2018<br>Publication | 1 %  |
| 7  | Nakamura, Shohei. "Generalized weighted Morrey spaces and classical operators : Generalized weighted Morrey spaces and classical operators", Mathematische Nachrichten, 2016.<br>Publication                                                 | 1 %  |
| 8  | Sun, H.. "Application of integral operator for regularized least-square regression", Mathematical and Computer Modelling, 200901<br>Publication                                                                                              | 1 %  |
| 9  | Submitted to Universitas Brawijaya<br>Student Paper                                                                                                                                                                                          | 1 %  |
| 10 | dokumen.pub<br>Internet Source                                                                                                                                                                                                               | 1 %  |
| 11 | www.researchgate.net<br>Internet Source                                                                                                                                                                                                      | <1 % |
| 12 | scitepress.org<br>Internet Source                                                                                                                                                                                                            | <1 % |

13 Yoshihiro Sawano, Satoko Sugano, Hitoshi Tanaka. "Orlicz–Morrey Spaces and Fractional Operators", Potential Analysis, 2011  
Publication  $<1\%$

---

14 repository.uin-malang.ac.id  
Internet Source  $<1\%$

---

15 Hendra Gunawan, Denny I. Hakim, Kevin M. Limanta, Al A. Masta. "Inclusion properties of generalized Morrey spaces", Mathematische Nachrichten, 2016  
Publication  $<1\%$

---

16 Mariano Giaquinta, Stefan Hildebrandt. "Calculus of Variations I", Springer Science and Business Media LLC, 2004  
Publication  $<1\%$

---

17 SANGHACK LEE. "TWO-STAGE CONTESTS WITH ADDITIVE CARRYOVERS\*", International Economic Journal, 2003  
Publication  $<1\%$

---

18 Vagif S. Guliyev, Kamala R. Rahimova. "Parabolic Fractional Maximal Operator in Modified Parabolic Morrey Spaces", Journal of Function Spaces and Applications, 2012  
Publication  $<1\%$

---

19 ccenet.org  
Internet Source  $<1\%$

---

21

"Differential and Integral Inequalities",  
Springer Science and Business Media LLC,  
2019

Publication

&lt;1 %

22

V. S. Guliyev. "Generalized Sobolev–Morrey  
estimates for hypoelliptic operators on  
homogeneous groups", Revista de la Real  
Academia de Ciencias Exactas, Físicas y  
Naturales. Serie A. Matemáticas, 2021

Publication

&lt;1 %

23

Vakhtang Kokilashvili, Alexander Meskhi,  
Humberto Rafeiro, Stefan Samko. "Integral  
Operators in Non-Standard Function Spaces",  
Springer Science and Business Media LLC,  
2016

Publication

&lt;1 %

24

Yanlong Shi, Xiangxing Tao. "SOME MULTI-  
SUBLINEAR OPERATORS ON GENERALIZED  
MORREY SPACES WITH NON-DOUBLING  
MEASURES", Journal of the Korean  
Mathematical Society, 2012

Publication

&lt;1 %

# revisi Bessel

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8