

Jensen m -convexity on set valued function.pdf

by Perpustakaan UM Surabaya

Submission date: 10-May-2024 01:19PM (UTC+0700)

Submission ID: 2375801799

File name: Jensen_m-convexity_on_set_valued_function.pdf (286.6K)

Word count: 3290

Character count: 14361



Jensen m-Convexity on Set-Valued Function

Eko Dedi Pramana*, Ratno Bagus Edy Wibowo, Mohamad Muslikh

13
¹Department of Mathematics, Faculty of Mathematics and Natural Sciences,
Universitas Brawijaya, Indonesia.
Email: eko.d.pramana@gmail.com.

ABSTRACT

5
This paper discusses the class of Jensen m-convex functions and their properties, leading to enhancements of these functions and Jensen's inequalities for set-valued functions. It provides several characterizations and explores various algebraic properties. Additionally, it introduces a discrete Jensen-type inclusion.

Keywords: Jensen m-convex; Jensen-type inclusion; Jensen m-convex set; Set-valued function.

INTRODUCTION

Convexity theory is a fundamental aspect of mathematical analysis with wide-ranging applications across various disciplines [1]. One significant development within this theory is the notion of m-convexity, a generalization of convexity introduced to extend the scope of traditional convex functions (See [2] [3] [4] [5]). In recent years, interest has grown significantly in studying m-convex functions and their properties, particularly in the context of set-valued functions (See [6] [7] [8]).

Jensen's inequality is a fundamental concept in convex analysis, offering crucial perspectives on the behavior of convex functions [9]. The extension of this inequality to set-valued functions is a dynamic research area, motivated by its potential applications in optimization, economics, statistics, and various other disciplines. Comprehending the attributes and features of set-valued functions within the framework of m-convexity is pivotal for tackling intricate optimization challenges and representing the uncertainties that are intrinsic to real-world scenarios [8].

Lara has provided the class of convexity in various fields such as operator theory, linear spaces, and functional analysis(see [10] [11] [12] [13] [14]). On the other hand, Lara delivered a strong version of Jensen m-convexity(See [15] [16]), which remains unexplored in certain areas, particularly in the concept of Set-Valued functions.

The primary goal of this research is to present the concept of a Jensen m-convex set-valued function and to elucidate specific properties of these functions, thereby generalizing the established concept of convexity for set-valued functions. These functions are defined on a nonempty Jensen m-convex subset of a real linear space, taking values in the set of nonempty subsets of another real linear space.

METHODS

The methodology employed in composing this research article involves a literature review of various related articles and books. Within this literature, both the definition

and properties of the m -convex function have been explored, as well as the characteristics of m -convexity in set-valued functions. Thus, utilizing the definitions and properties outlined in prior studies, this research investigates Jensen m -convexity in set-valued functions. Initially, the discussion focuses on Jensen m -convexity and m -convexity. ¹

Let Ω be a non-empty subset of $\mathbb{R}^n$. Ω is defined as a convex set if, for any x, y in Ω and any t in $[0,1]$, the combination $tx + (1-t)y$ also belongs to Ω . Geometrically, this implies that the line segment connecting any two points in Ω lies entirely within Ω . Interval is denoted I .

Definition 1 (m -Convex set, [6])

Let X be a real linear space and $m \in [0,1]$. A non-empty set $D \subseteq X$ is called m -convex set if for any $x, y \in D$ and $t \in [0,1]$ the point $tx + m(1-t)y \in D$.

Definition 2 (m -Convex function, [2] [3] [10] [12])

Let $m \in [0,1]$. A function $f: [0, b] \rightarrow \mathbb{R} (b > 0)$ is said to be m -convex in the interval $[0, b]$, if for any $x, y \in [0, b]$ and $t \in [0,1]$ we have

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y). \quad (1)$$

⁶ The concept of an m -convex function can be interpreted geometrically. a function $f: [0, b] \rightarrow \mathbb{R}$ is considered m -convex if, for any $x, y \in [0, b]$ with $x \leq y$, the line segment connecting the points $(x, f(x))$ and $(y, mf(y))$ lies above the graph of f on the interval $[x, my]$. ⁶

⁷ The foundations of convex function theory are attributed to Jensen. It is also noted that the original definition of a real-valued convex function, as established by Jensen, is what is now commonly referred to as a mid-convex function or Jensen convex. A Jensen convex function is a function $f: I \rightarrow \mathbb{R}$ that satisfies the following inequality ¹

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x)+f(y)}{2}, \quad (2)$$

for all $x, y \in I$.

Definition 3 (Jensen m -convex function, [15])

Let $m \in (0,1]$. A function $f: [0, +\infty) \rightarrow \mathbb{R}$ is called Jensen m -convex function in the interval $[0, +\infty)$ if satisfies the inequality

$$f\left(\frac{x+y}{c_m}\right) \leq \frac{f(x)+f(y)}{c_m}, \quad (3)$$

for all $x, y \in [0, +\infty)$.

¹² The concept of a Jensen m -convex function pertains to each m within the interval $(0,1]$, for a specific value t in the open interval $(0,1)$, which is denoted with respect to m as t_m . Given any $(x, y) \in [0, b] \times [0, b]$, $t_mx + m(1-t_m)y$ can be represented as

$$t_mx + m(1-t_m)y = \frac{x+y}{c_m}, \quad (4)$$

where c_m is a constant that depends only on m . By rewriting Equation (4) as

$$\left(t_m - \frac{1}{c_m}\right)x + \left(m(1-t_m) - \frac{1}{c_m}\right)y = 0,$$

and solving for c_m and t_m we obtain

$$c_m = 1 + \frac{1}{m}, \quad \text{and} \quad t_m = \frac{m}{m+1}. \quad (5)$$

Definition 4 (*starshaped function*, [15])

A function $f: I \rightarrow \mathbb{R}$ that satisfies the inequality

$$f(tx) \leq tf(x), \quad (6)$$

for all $x \in I$ and $t \in [0,1]$ will be called *starshaped function* in the interval I .

Since Jensen m -convex functions exhibit properties as outlined in (5), the concept of *starshaped functions* are utilized to produce algebraic properties within Jensen m -convex functions.

Definition 5 (m -Convex set-valued function, [6])

Let $m \in [0,1]$. Let D be any non-empty m -convex subset of X . A set-valued function $F: D \rightarrow n(Y)$ is called m -convex if satisfies the inclusion

$$tF(x) + m(1-t)F(y) \subseteq F(tx + m(1-t)y), \quad (7)$$

for all $x, y \in D$ and $t \in [0,1]$.

RESULTS AND DISCUSSION

In this section, we present some results on Jensen m -convexity for set-valued functions. Our focus extends the concept of Jensen m -convexity to set-valued functions. Let X and Y denote real linear spaces, $n(Y)$ represent the family of all non-empty subsets of Y , and let m and t_m be fixed real numbers in the interval $(0,1]$.

Definition 6 Let X be a real linear space and $m \in (0,1]$. A non-empty set $D_m \subseteq X$ is called Jensen m -convex if for any $x, y \in D_m$ the point $\frac{x+y}{c_m} \in D_m$.

Example 1 Set $A = \{x | 0 \leq x \leq b; x, y \in \mathbb{Q}\}$ is Jensen m -convex set for rational number $m \in (0,1]$.

Definition 7 Let $m \in (0,1]$ and D_m be a Jensen m -convex set. Set-valued function $F: D_m \rightarrow n(Y)$ is said to be Jensen m -convex if satisfies the inclusion

$$\frac{F(x) + F(y)}{c_m} \subseteq F\left(\frac{x+y}{c_m}\right), \quad (8)$$

for all $x, y \in D_m$ with c_m is a constant value depend on m .

The distinction between Jensen m -convex and m -convex functions lies in the role of m within the inequality. In Definition 2 and Definition 5, the value of t is arbitrary within the interval $[0,1]$, whereas for Jensen m -convex functions, the value of t_m is dependent on m . From Inclusion (8) by taking value $x = y$, we have

$$\frac{2F(x)}{c_m} \subseteq F\left(\frac{2x}{c_m}\right),$$

given that $\lambda_m = \frac{2}{c_m}$, we obtain

$$\lambda_m F(x) \subseteq F(\lambda_m x). \quad (9)$$

Proposition 1 Let $m \in (0,1]$. If $F: D_m \rightarrow n(Y)$ is a Jensen m -convex set-valued function and starshaped function, then F satisfies the inclusion

$$\frac{1}{c_m} [tF(c_m x) + (1-t)F(c_m y)] \subseteq F(tx + (1-t)y), \quad (10)$$

for any $x, y \in D_m$ and $t \in (0,1)$.

Proof.

Let $x, y \in D_m$ and $t \in (0,1)$, then we have

$$\begin{aligned} \frac{1}{c_m} [tF(c_mx) + (1-t)F(c_my)] &\subseteq \frac{1}{c_m} [F(c_mtx) + F(c_m(1-t)y)] \\ &\subseteq F\left(\frac{c_mtx + c_m(1-t)y}{c_m}\right) \\ &= F(tx + (1-t)y) \end{aligned}$$

Proposition 2 Let $m \in (0,1]$. If $F: D_m \rightarrow n(Y)$ is a Jensen m -convex set-valued function and starshaped function, then for all positive real number $x_1, x_2, y_1, y_2 \in D_m$ we have

$$\frac{1}{c_m} \left[y_1 F\left(\frac{c_mx_1}{y_1}\right) + y_2 F\left(\frac{c_mx_2}{y_2}\right) \right] \subseteq (y_1 + y_2) F\left(\frac{x_1 + x_2}{y_1 + y_2}\right). \quad (11)$$

Proof.

Let F Jensen m -convex set-valued function and starshaped, then for all real positive $x_1, x_2, y_1, y_2 \in D_m$ we have

$$\begin{aligned} &\frac{1}{c_m} \left[y_1 F\left(\frac{c_mx_1}{y_1}\right) + y_2 F\left(\frac{c_mx_2}{y_2}\right) \right] \\ &= \frac{y_1 + y_2}{c_m} \left[\frac{y_1}{y_1 + y_2} F\left(\frac{c_mx_1}{y_1}\right) + \frac{y_2}{y_1 + y_2} F\left(\frac{c_mx_2}{y_2}\right) \right] \\ &= \frac{y_1 + y_2}{c_m} \left[\frac{y_1}{y_1 + y_2} F\left(\frac{c_mx_1}{y_1}\right) + \left(1 - \frac{y_1}{y_1 + y_2}\right) F\left(\frac{c_mx_2}{y_2}\right) \right] \\ &\subseteq (y_1 + y_2) F\left(\frac{x_1 + x_2}{y_1 + y_2}\right) \end{aligned}$$

Proposition 3 Let $m \in (0,1]$. If $F: D_m \rightarrow n(Y)$ Jensen m -convex, then image of F for any subset Jensen m -convex of D_m are Jensen m -convex set of linear space Y .

Proof.

Let A be a non-empty Jensen m -convex subset of D_m , and let a, b belong to $F(A)$, where $F(A)$ represents the union of F over $x \in A$. Then, by taking $a \in F(x)$ and $b \in F(y)$ for some $x, y \in A$, we have

$$\frac{a + b}{c_m} \in \frac{F(x) + F(y)}{c_m} \subseteq F\left(\frac{(x + y)}{c_m}\right).$$

Since A is Jensen m -convex, $\frac{x+y}{c_m} \in A$, implying that $\frac{a+b}{c_m} \in F(A)$. Therefore, $F(A)$ is a Jensen m -convex function on Y .

Theorem 1 Let $F: D_m \subseteq X \rightarrow n(Y)$ be a set-valued function. F is said to be Jensen m -convex function if and only if its graph, denoted by $Gr_m(F) := \{(x, y) \in X \times Y \mid x \in D_m, y \in F(x)\}$, is Jensen m -convex set in $X \times Y$.

Proof.

($\Rightarrow$) Let $m \in (0,1]$. For any $(x_1, y_1), (x_2, y_2) \in Gr_m$ such that $y_1 \in F(x_1)$ and $y_2 \in F(x_2)$ and satisfies $\frac{y_1 + y_2}{c_m} \in \frac{F(x_1) + F(x_2)}{c_m}$. Since, F is Jensen m -convex we have $\left(\frac{x_1 + x_2}{c_m}, \frac{y_1 + y_2}{c_m}\right) \in Gr_m$ that implies,

$$\frac{(x_1, y_1) + (x_2, y_2)}{c_m} \in Gr_m(F).$$

($\Leftarrow$) Let $m \in (0,1]$. For any $x, y \in D_m$, select $y \in \frac{F(x_1)+F(x_2)}{c_m}$ such that y can be expressed as $\frac{y_1+y_2}{c_m}$ where $y_1 \in F(x_1)$ and $y_2 \in F(x_2)$. Suppose Gr_m is a Jensen m -convex set over $X \times Y$ with respect to m . Then for $(x_1, y_1), (x_2, y_2) \in Gr_m$, we have $\frac{(x_1, y_1) + (x_2, y_2)}{c_m} \in Gr_m(F)$, that implies

$$\left(\frac{x_1 + x_2}{c_m}, \frac{y_1 + y_2}{c_m} \right) \in Gr_m.$$

Hence, we have $y = \frac{y_1+y_2}{c_m} \in F\left(\frac{x_1+x_2}{c_m}\right)$. Given that y is constructed from any $x_1, x_2 \in D_m$, it follows that F is Jensen m -convex.

Definition 8 Let $F_1, F_2: D_m \subseteq X \rightarrow n(Y)$ be a set-valued function, we have

- 1) The union function of F_1 and F_2 is a set-valued function $F_1 \cup F_2: D_m \rightarrow n(Y)$ which defined by $(F_1 \cup F_2)(x) = F_1(x) \cup F_2(x)$ for all $x \in D_m$.
- 2) The intersection function of F_1 and F_2 is a set-valued function $F_1 \cap F_2: D_m \rightarrow n(Y)$ which defined by $(F_1 \cap F_2)(x) = F_1(x) \cap F_2(x)$ for all $x \in D_m$.
- 3) The sum function of F_1 and F_2 is a set-valued function $(F_1 + F_2)(x) = F_1(x) + F_2(x)$ for all $x \in D_m$.

Definition 9 Let X, Y, Z be linear spaces and let D_m be a subset of X . Then we have

- 1) If $F_1: D_m \rightarrow n(Y)$ and $F_2: D_m \rightarrow n(Z)$ are set-valued functions. Then the Cartesian product of F_1 and F_2 is set-valued function $(F_1 \times F_2)(x): D_m \rightarrow n(Y) \times n(Z)$ which defined by $(F_1 \times F_2)(x) = F_1(x) \times F_2(x)$ for all $x \in D_m$.
- 2) If $F_1: D_m \rightarrow n(Y)$ and $F_2: n(Y) \rightarrow n(Z)$ are set-valued function. Then the composition function of F_1 and F_2 is the set-valued function $F_2 \circ F_1: D_m \rightarrow n(Z)$ given by $(F_2 \circ F_1)(x) = F_2(F_1(x)) = \bigcup_{y \in F_1(x)} F_2(y)$ for all $x \in D_m$.

Proposition 4 Let $m \in (0,1]$. If $F_1, F_2: D_m \rightarrow n(Y)$ are Jensen m -convex set-valued functions which satisfies

$$F_1(x) \subseteq F_2(x) \text{ (or } F_2(x) \subseteq F_1(x)),$$

for each $x \in D_m$. Then the union function of F_1 and F_2 is Jensen m -convex set-valued function.

Proof Let $m \in (0,1]$ and $x_1, x_2 \in D_m$. Without loss of generalization, let $F_1(x) \subseteq F_2(x)$. Then we have

$$\begin{aligned} \frac{(F_1 \cup F_2)(x_1) + (F_1 \cup F_2)(x_2)}{c_m} &= \frac{F_1(x_1) \cup F_2(x_1) + F_1(x_2) \cup F_2(x_2)}{c_m} \\ &= \frac{F_2(x_1) + F_2(x_2)}{c_m} \\ &\subseteq F_2\left(\frac{x_1 + x_2}{c_m}\right) \\ &= F_1\left(\frac{x_1 + x_2}{c_m}\right) \cup F_2\left(\frac{x_1 + x_2}{c_m}\right) \\ &= (F_1 \cup F_2)\left(\frac{x_1 + x_2}{c_m}\right) \end{aligned}$$

Furthermore, $(x, y) \in Gr_m(F_1 \cup F_2)$ if and only if $y \in (F_1 \cup F_2)(x) = F_1(x) \cup F_2(x)$ which

is equivalent with

$$Gr_m(F_1 \cup F_2) = Gr_m(F_1) \cup Gr_m(F_2). \quad (12)$$

Proposition 5 Let $m \in (0,1]$. If $F_1, F_2: D_m \rightarrow n(Y)$ are Jensen m -convex set-valued functions. Then the intersection function of F_1 and F_2 is Jensen m -convex set-valued function.

Proof Since F_1 and F_2 are Jensen m -convex set-valued function, we have $Gr_m(F_1)$ and $Gr_m(F_2)$ are Jensen m -convex and

$$\begin{aligned} Gr_m(F_1) \cap Gr_m(F_2) &= \{(x, y) | x \in D_m(F_1) \cap D_m(F_2), y \in F_1(x) \cap F_2(x)\} \\ &= \{(x, y) | x \in D_m(F_1 \cap F_2), y \in (F_1 \cap F_2)(x)\} \\ &= Gr_m(F_1 \cap F_2). \end{aligned}$$

We can conclude that $Gr_m(F_1 \cap F_2)$ is Jensen m -convex. Then by Theorem 1, $F_1 \cap F_2$ is Jensen m -convex set-valued function.

Proposition 6 Let $m \in (0,1]$. If $F_1, F_2: D_m \rightarrow n(Y)$ are Jensen m -convex set-valued functions. Then the sum function of F_1 and F_2 is Jensen m -convex set-valued function.

Proof Let $x, y \in D_m$, since F_1 and F_2 Jensen m -convex set-valued function we have

$$\begin{aligned} \frac{(F_1 + F_2)(x) + (F_1 + F_2)(y)}{c_m} &= \frac{F_1(x) + F_2(x) + F_1(y) + F_2(y)}{c_m} \\ &= \frac{F_1(x) + F_1(y)}{c_m} + \frac{F_2(x) + F_2(y)}{c_m} \\ &\subseteq F_1\left(\frac{x+y}{c_m}\right) + F_2\left(\frac{x+y}{c_m}\right) \\ &= (F_1 + F_2)\left(\frac{x+y}{c_m}\right). \end{aligned}$$

Proposition 7 Let $m \in (0,1]$. If $F_1, F_2: D_m \rightarrow n(Y)$ are Jensen m -convex set-valued functions. Then the Cartesian product of F_1 and F_2 is Jensen m -convex set-valued function.

Proof Let $x, y \in D_m$, since F_1 and F_2 Jensen m -convex set-valued function we have

$$\begin{aligned} \frac{(F_1 \times F_2)(x) + (F_1 \times F_2)(y)}{c_m} &= \frac{F_1(x) \times F_2(x) + F_1(y) \times F_2(y)}{c_m} \\ &= \frac{(F_1(x) + F_1(y)) \times (F_2(x) + F_2(y))}{c_m} \\ &= \frac{F_1(x) + F_1(y)}{c_m} \times \frac{F_2(x) + F_2(y)}{c_m} \\ &\subseteq F_1\left(\frac{x+y}{c_m}\right) \times F_2\left(\frac{x+y}{c_m}\right) = (F_1 \times F_2)\left(\frac{x+y}{c_m}\right). \end{aligned}$$

Proposition 8 Let $m \in (0,1]$. If $F_1: D_m \rightarrow n(Y)$ and $F_2: n(Y) \rightarrow n(Z)$ are Jensen m -convex set-valued functions. Then the composition function $(F_2 \circ F_1)$ is Jensen m -convex set-valued function.

Proof Let $x, y \in D_m$, since F_1 and F_2 Jensen m -convex set-valued function we have

$$\begin{aligned}
\frac{(F_2 \circ F_1)(x) + (F_2 \circ F_1)(y)}{c_m} &= \frac{F_2(F_1(x)) + F_2(F_1(y))}{c_m} \\
&\subseteq F_2\left(\frac{F_1(x) + F_1(y)}{c_m}\right) \\
&\subseteq F_2\left(F_1\left(\frac{x+y}{c_m}\right)\right) \\
&= (F_1 \circ F_2)\left(\frac{x+y}{c_m}\right).
\end{aligned}$$

■

Theorem 2 Let $m \in (0,1]$ and a function $F: D_m \rightarrow n(Y)$ be a Jensen m -convex set-valued function and starshaped. Then for $n \geq 2$ and $A_1, A_2, \dots, A_n$ are non-empty subset of D_m satisfies the inclusion,

$$\frac{1}{n} \left[\frac{1}{c_m^{n-1}} F(c_m^{n-1} A_1) + \sum_{k=1}^{n-1} \frac{F(c_m^{n-k} A_{k+1})}{c_m^{n-k}} \right] \subseteq F\left(\frac{1}{n} \sum_{k=1}^n A_k\right). \quad (13)$$

Proof Since F is starshaped, for all n we have

$$\frac{1}{n} F\left(\sum_{k=1}^n A_k\right) \subseteq F\left(\frac{1}{n} \sum_{k=1}^n A_k\right).$$

Therefore, to establish the argument, it suffices to demonstrate the following inclusions,

$$\frac{1}{c_m^{n-1}} F(c_m^{n-1} A_1) + \sum_{k=1}^{n-1} \frac{F(c_m^{n-k} A_{k+1})}{c_m^{n-k}} \subseteq F\left(\frac{1}{n} \sum_{k=1}^n A_k\right).$$

For $n = 2$, by applying Proposition 1 and selecting $t = \frac{1}{2}$, we obtain

$$\frac{1}{2} \left[\frac{F(c_m A_1)}{c_m} + \frac{F(c_m A_2)}{c_m} \right] \subseteq F\left(\frac{1}{2} (A_1 + A_2)\right).$$

Assume that Inclusion (13) holds for n , now for $n + 1$

$$\begin{aligned}
&\frac{1}{c_m^n} F(c_m^n A_1) + \sum_{k=1}^n \frac{F(c_m^{n+1-k} A_{k+1})}{c_m^{n+1-k}} \\
&= \frac{1}{c_m} \left[\frac{1}{c_m^{n-1}} F(c_m^{n-1} c_m A_1) + \sum_{k=1}^{n-1} \frac{F(c_m^{n-k} c_m A_{k+1})}{c_m^{n-k}} \right] + \frac{1}{c_m} F(c_m A_{n+1}) \\
&\subseteq \frac{1}{c_m} \left[F\left(\sum_{k=1}^n c_m A_k\right) + F(c_m A_{n+1}) \right] \\
&\subseteq F\left(\frac{\sum_{k=1}^n c_m A_k + c_m A_{n+1}}{c_m}\right) = F\left(\sum_{k=1}^{n+1} A_k\right).
\end{aligned}$$

■

Corollary 1 Let $m \in (0,1]$ and a function $F: D_m \rightarrow n(Y)$ be a Jensen m -convex set-valued function and starshaped. Then for $n \geq 2$ and $x_1, x_2, \dots, x_n \in D_m$ satisfies the inclusion,

$$\frac{1}{n} \left[\frac{1}{c_m^{n-1}} F(c_m^{n-1} x_1) + \sum_{k=1}^{n-1} \frac{F(c_m^{n-k} x_{k+1})}{c_m^{n-k}} \right] \subseteq F\left(\frac{1}{n} \sum_{k=1}^n x_k\right). \quad (14)$$

CONCLUSIONS

Jensen m -convex set-valued function attain the same some properties from m -convexity set-valued function. We present example of Jensen m -convex set and the algebraic properties. Specifically, we establish the discrete Jensen-type inclusion of Jensen m -convex set-valued function.

ACKNOWLEDGMENTS

The author extends sincere thanks for the constructive comments from the referee, which have greatly improved the manuscript. Furthermore, the author appreciates the partial financial support from LPDP Scholarship.

REFERENCES

- [1] J. R. Džurlewick, "A short history of Convexity," *Differential Geometry - Dynamical Systems*, vol. 11, pp. 112-129, 2009.
- [2] S. S. Dragomir and G. Toader, "Some Inequalities for m -Convex Functions.," *Studia Univ. Babeş-Bolyai, Math*, vol. 38, no. 1, pp. 21-28, 1993.
- [3] S. S. Dragomir, "On some new inequalities of Hermite-Hadamard type for m -convex functions," *Tamkang J. of Math*, vol. 33, no. 1, pp. 45-55, 2002.
- [4] L. Hormander, *Notions of Convexity*, vol. 207, Basel: Birkhäuser, 1994.
- [5] M. K. Bakula, M. E. Ozdemir and J. Pecaric, "Hadamard Type Inequalities For m -Convex and (α, m) -Convex," *Journal of Inequalities in Pure and Applied Mathematics*, vol. 9, no. 4, pp. 96-108, 2008.
- [6] T. Lara, N. Marentes, R. Quintero and E. Rosales, "On m -Convexity of Set-Valued Functions," *Advances in Operator Theory*, vol. 4, no. 1, pp. 1-16, 2019.
- [7] T. Lara, N. Marentes, R. Quintero and E. Rosales, "Strong m -Convexity of Set-Valued Functions," *Annales Mathematicae Silesianae*, vol. 37, pp. 1-13, 2023.
- [8] D. Zhang, C. Guo, D. Chen and G. Wang, "Jensen's inequalities for set-valued and fuzzy set-valued functions," *Fuzzy Sets and Systems*, vol. 404, pp. 178-204, 2021.
- [9] S. I. Bradanović, "Improvements of Jensen's inequality and its converse for strongly convex functions with applications to strongly f -divergences," *Journal of Mathematical Analysis and Applications*, vol. 531, no. 2, p. 127866, 2024.
- [10] T. Lara, E. Rosales and J. Sanchez, "New properties of m -convex functions," *International Journal of Mathematical Analysis*, vol. 9, no. 15, pp. 735-742, 2015.
- [11] L. Berbesi, T. Lara, P. Pena and E. Rosales, "On operator m -convex functions in hilbert space," *UPI Journal of Mathematics and Biostatistics*, vol. 1, no. 2, pp. 1-12, 2018.
- [12] T. Lara, N. Marentes, Z. Pales and R. Quintero, "On m -convexity on real linear spaces," *UPI Journal of Mathematics and Biostatistics*, vol. 1, no. 2, pp. 1-16, 2018.
- [13] T. Lara, N. Marentes, R. Quintero and E. Rosales, "On m -concave functions on real linear spaces," *Boletín de la Asociación Matemática Venezolana*, vol. 23, no. 2, pp. 131-137, 2016.
- [14] T. Lara, N. Marentes, R. Quintero and E. Rosales, "Stability of m -jensen functional equations," *Journal of Advances in Mathematics and Computer Science*, vol. 24, no. 1, pp. 1-12, 2017.
- [15] T. Lara, R. Quintero, E. Rosales and J. Sanchez, "On Strongly Jensen m -Convex Function," *Pure Mathematical Sciences*, vol. 6, no. 1, pp. 87-94, 2017.
- [16] T. Lara, R. Quintero, E. Rosales and J. Sanchez, "On a generalization of the class of Jensen convex functions," *Aequationes Mathematicae*, vol. 90, 2016.

Jensen m-convexity on set valued function.pdf

ORIGINALITY REPORT

14%

SIMILARITY INDEX

%

INTERNET SOURCES

13%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | Basic Modern Algebra with Applications, 2014.
Publication | 1 % |
| 2 | Bot, R.I.. "An alternative formulation for a new closed cone constraint qualification", Nonlinear Analysis, 20060315
Publication | 1 % |
| 3 | Nonconvex Optimization and Its Applications, 2005.
Publication | 1 % |
| 4 | Friedrich Knop. "Multiplicities and Dimensions in Enveloping Tensor Categories", International Mathematics Research Notices, 2024
Publication | 1 % |
| 5 | Teodoro Lara, Roy Quintero, Edgar Rosales, José Luis Sánchez. "On a generalization of the class of Jensen convex functions", Aequationes mathematicae, 2016
Publication | 1 % |

6	Teodoro Lara, Edgar Rosales, Jose L. Sanchez. "New properties of m-convex functions", International Journal of Mathematical Analysis, 2015 Publication	1 %
7	Teodoro Lara, Nelson Merentes, Roy Quintero, Edgar Rosales. "Properties of Jensen m-convex functions", International Journal of Mathematical Analysis, 2016 Publication	1 %
8	Teodoro Lara, Nelson Merentes, Roy Quintero, Edgar Rosales. " Strong -Convexity of Set-Valued Functions ", Annales Mathematicae Silesianae, 2023 Publication	1 %
9	Submitted to Universitas Pendidikan Indonesia Student Paper	1 %
10	Jorge Cortes. "Discontinuous dynamical systems", IEEE Control Systems Magazine, 6/2008 Publication	1 %
11	Submitted to University of Huddersfield Student Paper	<1 %
12	Teodoro Lara, Roy Quintero, Edgar Rosales, Jose L. Sanchez. "On strongly Jensen m-	<1 %

convex functions", Pure Mathematical Sciences, 2017

Publication

13

Submitted to Universitas Brawijaya

Student Paper

<1 %

14

Hamid Khodaei. "Selections of generalized convex set-valued functions satisfying some inclusions", Journal of Mathematical Analysis and Applications, 2019

Publication

<1 %

15

J. Aczél, F. Stehling. "The Twenty-seventh International Symposium on Functional Equations, August 14–24, 1989, Bielsko-BiałKatowice—Kraków, Poland", Aequationes Mathematicae, 1990

Publication

<1 %

16

M. D. Fajardo. "Set-Valued Evenly Convex Functions: Characterizations and C-Conjugacy", Set-Valued and Variational Analysis, 2022

Publication

<1 %

17

Rockafellar. "Max and Min", Grundlehren der mathematischen Wissenschaften, 1998

Publication

<1 %

18

Slavica Ivelić Bradanović. "Improvements of Jensen's inequality and its converse for strongly convex functions with applications to

<1 %

strongly f-divergences", Journal of
Mathematical Analysis and Applications, 2024

Publication

19

Thales Sarinho Galvão Santos de Souza.
"Open Coloring Axiom e aplicações de
colorações", Universidade de São Paulo.
Agência de Bibliotecas e Coleções Digitais,
2023

Publication

<1 %

20

Erhan Set, Maryam Sardari, Muhamet Emin
Ozdemir, Jamal Rooin. "On Generalizations of
the Hadamard Inequality for (α, m) -Convex
Functions", Kyungpook mathematical journal,
2012

Publication

<1 %

21

Erhan Set. "On Hadamard-Type Inequalities
Involving Several Kinds of Convexity", Journal
of Inequalities and Applications, 2010

Publication

<1 %

22

Subir Bose, George Deltas. "Exclusive Versus
Non-exclusive Dealing in Auctions with
Resale", Economic Theory, 2006

Publication

<1 %

23

X. M. Yang, D. Li, S. Y. Wang. "Near-
Subconvexlikeness in Vector Optimization
with Set-Valued Functions", Journal of
Optimization Theory and Applications, 2001

Publication

<1 %

24

V.F. Demyanov. " Exhausters af a positively homogeneous function ", Optimization, 1999

Publication

<1 %

25

"Vector Optimization", Springer Nature, 2005

Publication

<1 %

26

Submitted to Queen's University of Belfast

Student Paper

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On